

WEST Search History

Inventor
search

DATE: Tuesday, September 11, 2007

<u>Hide?</u>	<u>Set Name</u>	<u>Query</u>	<u>Hit Count</u>
		<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>	
<input type="checkbox"/>	L4	L3 and aurora-A	2
<input type="checkbox"/>	L3	(martin anne)[IN]	13
<input type="checkbox"/>	L2	L1 and aurora-A	2
<input type="checkbox"/>	L1	(prigent claudie)[IN]	6

END OF SEARCH HISTORY

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTALAB1643

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	MAY 01	New CAS web site launched
NEWS	3	MAY 08	CA/CAPplus Indian patent publication number format defined
NEWS	4	MAY 14	RDISCLOSURE on STN Easy enhanced with new search and display fields
NEWS	5	MAY 21	BIOSIS reloaded and enhanced with archival data
NEWS	6	MAY 21	TOXCENTER enhanced with BIOSIS reload
NEWS	7	MAY 21	CA/CAPplus enhanced with additional kind codes for German patents
NEWS	8	MAY 22	CA/CAPplus enhanced with IPC reclassification in Japanese patents
NEWS	9	JUN 27	CA/CAPplus enhanced with pre-1967 CAS Registry Numbers
NEWS	10	JUN 29	STN Viewer now available
NEWS	11	JUN 29	STN Express, Version 8.2, now available
NEWS	12	JUL 02	LEMBASE coverage updated
NEWS	13	JUL 02	LMEDLINE coverage updated
NEWS	14	JUL 02	SCISEARCH enhanced with complete author names
NEWS	15	JUL 02	CHEMCATS accession numbers revised
NEWS	16	JUL 02	CA/CAPplus enhanced with utility model patents from China
NEWS	17	JUL 16	CAPplus enhanced with French and German abstracts
NEWS	18	JUL 18	CA/CAPplus patent coverage enhanced
NEWS	19	JUL 26	USPATFULL/USPAT2 enhanced with IPC reclassification
NEWS	20	JUL 30	USGENE now available on STN
NEWS	21	AUG 06	CAS REGISTRY enhanced with new experimental property tags
NEWS	22	AUG 06	BEILSTEIN updated with new compounds
NEWS	23	AUG 06	FSTA enhanced with new thesaurus edition
NEWS	24	AUG 13	CA/CAPplus enhanced with additional kind codes for granted patents
NEWS	25	AUG 20	CA/CAPplus enhanced with CAS indexing in pre-1907 records
NEWS	26	AUG 27	Full-text patent databases enhanced with predefined patent family display formats from INPADOCDB
NEWS	27	AUG 27	USPATOLD now available on STN
NEWS	28	AUG 28	CAS REGISTRY enhanced with additional experimental spectral property data
NEWS	29	SEP 07	STN AnaVist, Version 2.0, now available with Derwent World Patents Index
NEWS EXPRESS	05	SEPTEMBER 2007:	CURRENT WINDOWS VERSION IS V8.2, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 05 SEPTEMBER 2007.
NEWS HOURS			STN Operating Hours Plus Help Desk Availability
NEWS LOGIN			Welcome Banner and News Items
NEWS IPC8			For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 09:28:02 ON 11 SEP 2007

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'CAPLUS' ENTERED AT 09:28:15 ON 11 SEP 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 11 Sep 2007 VOL 147 ISS 12

FILE LAST UPDATED: 10 Sep 2007 (20070910/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s (Aurora-2) or (Aur-2) or (STK-15) or (AIK) or (ARK1) or (AurA) or (AURA)

4145 AURORA

495 AURORAS

4226 AURORA

(AURORA OR AURORAS)

9285941 2

113 AURORA-2

(AURORA(W)2)

698 AUR

1 AURS

699 AUR

(AUR OR AURS)

9285941 2

12 AUR-2

(AUR(W)2)

424 STK

31 STKS

452 STK

(STK OR STKS)

1750137 15

4 STK-15

(STK(W)15)

130 AIK

1 AIKS

130 AIK
 (AIK OR AIKS)
 184 ARK1
 586 AURA
 7 AURAS
 593 AURA
 (AURA OR AURAS)
 586 AURA
 7 AURAS
 593 AURA

(AURA OR AURAS)
 L1 1027 (AURORA-2) OR (AUR-2) OR (STK-15) OR (AIK) OR (ARK1) OR (AURA)
 OR (AURA)

=> s L1 and (monoclonal)
 148722 MONOCLONAL
 543 MONOCLONALS
 148789 MONOCLONAL
 (MONOCLONAL OR MONOCLONALS)

L2 8 L1 AND (MONOCLONAL)

=> duplicate remove L2
 PROCESSING COMPLETED FOR L2
 L3 8 DUPLICATE REMOVE L2 (0 DUPLICATES REMOVED)

=> d L3 bib abs 1-8

L3 ANSWER 1 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN
 AN 2006:1158010 CAPLUS
 DN 145:469962
 TI Monoclonal anti-MCM2 protein antibodies for diagnosis,
 prevention and treatment of cervical diseases
 IN Fischer, Timothy J.; Malinowski, Douglas P.; Taylor, Adriann J.
 PA Tripath Imaging, Inc., USA
 SO PCT Int. Appl., 59pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2006116442	A2	20061102	WO 2006-US15706	20060426
	WO 2006116442	A3	20070705		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA				
	US 2006252106	A1	20061109	US 2006-410272	20060424
	US 2007117167	A1	20070524	US 2006-643277	20061221
PRAI	US 2005-675305P	P	20050427		
	US 2005-718082P	P	20050916		
	US 2006-410272	A3	20060424		

AB The invention provides a novel class of compds., pharmaceutical compns. comprising such compds. and methods of using such compds. to treat or prevent diseases or disorders associated with abnormal or deregulated kinase activity, particularly diseases or disorders that involve abnormal

activation of Aik, AbI, BRK, BIK, BMX, CSK, c-Src, c-Raf, EGFR, Fes, FGFR3, Fms, Fyn, IGF-IR, IR, IKK α , IKK β , JAK2, JAK3, KDR, Lck, Met, p70S6k, Ros, Rsk1, SAPK2 α , SAPK2 β , SAPK3, SIK, Tie2, TrkB and/or WNK3 kinases. The disease is a early stage HPV infection or cervical disease such as cervical carcinoma and mild dysplasia. The antibodies are specific to epitopes of MCM2 protein (Minichromosome maintenance protein 2).

L3 ANSWER 2 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN
 AN 2003:757393 CAPLUS
 DN 139:271089
 TI Phosphoinositide 3-kinase mediated inhibition of GPCRs
 IN Rockman, Howard A.; Naga, Prasad Sathyamangla V.; Laporte, Stephane A.; Barak, Larry S.; Caron, Marc G.
 PA USA
 SO U.S. Pat. Appl. Publ., 71 pp.
 CODEN: USXXCO

DT Patent
 LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2003182669	A1	20030925	US 2002-101235	20020319
	WO 2003088924	A2	20031030	WO 2003-US8208	20030318
	WO 2003088924	A3	20051229		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2003253586	A1	20031103	AU 2003-253586	20030318
	US 2006026702	A1	20060202	US 2004-902137	20040730
PRAI	US 2002-101235	A	20020319		
	WO 2003-US8208	W	20030318		

AB The present invention relates to compds. that alter G protein-coupled receptor (GPCR) internalization and new methods for their identification. Compds. of this invention include modified phosphoinositide 3-kinase (PI3K), modified HEAT domain, modified β -adrenergic receptor kinase 1 (β ARK1), as well as other peptides or small mols. that alter GPCR internalization. The present invention also includes the use of such compds. to treat GPCR-related diseases, such as cardiovascular disease, heart failure, asthma, nephrogenic diabetes insipidus, or hypertension.

L3 ANSWER 3 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN
 AN 2002:488124 CAPLUS
 DN 137:59517
 TI Human AURORA-1 and AURORA-2 kinases, cDNA and amino acid sequences, and recombinant production
 IN Plowman, Gregory; Mossie, Kevin
 PA Sugan, Inc., USA
 SO U.S. Pat. Appl. Publ., 43 pp., Cont.-in-part of U.S. Ser. No. 5,268, abandoned.
 CODEN: USXXCO

DT Patent
 LA English

FAN.CNT 4

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE

PI	US 2002081578	A1	20020627	US 1998-12135	19980122
	US 6716575	B2	20040406		
	CN 1205740	A	19990120	CN 1996-199101	19961125
	US 5962312	A	19991005	US 1996-755728	19961125
	EP 1655369	A1	20060510	EP 2005-23434	19961125
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
	CA 2318352	A1	19990729	CA 1999-2318352	19990121
	WO 9937788	A2	19990729	WO 1999-US1283	19990121
	WO 9937788	A3	19990930		
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
	RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	AU 9925605	A	19990809	AU 1999-25605	19990121
	EP 1051500	A2	20001115	EP 1999-905450	19990121
	EP 1051500	B1	20050817		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
	JP 2002508937	T	20020326	JP 2000-528695	19990121
	AT 302278	T	20050915	AT 1999-905450	19990121
	ES 2247783	T3	20060301	ES 1999-905450	19990121
	US 6207401	B1	20010327	US 1999-283011	19990331
	US 2005002938	A1	20050106	US 2001-784332	20010216
	US 6841579	B2	20050111		
	US 2004265852	A1	20041230	US 2003-734126	20031215
	US 7119174	B2	20061010		
PRAI	US 1995-8809P	P	19951218		
	US 1996-23943P	P	19960814		
	US 1996-755728	A2	19961125		
	US 1998-5268	B2	19980109		
	EP 1996-940870	A3	19961125		
	US 1998-12135	A	19980122		
	WO 1999-US1283	W	19990121		
	US 1999-283011	A3	19990331		
	US 2001-784332	A3	20010216		

AB The invention provides protein and cDNA sequences for human AURORA-1 (AUR1) and/or AURORA-2 (AUR2), which are members of serine/threonine kinase family containing short N-terminal extensions. AUR1 mRNA has been shown to be broadly expressed in rapidly dividing cells, derived from both normal and tumor tissues. AUR2 mRNA, however, has been shown to be expressed in a more restricted pattern being low or absent in most normal tissues and abundant in only a subset of tumor-derived cell lines. The invention also demonstrated that AUR1 and AUR2 kinases were able to phosphorylate myelin basic protein. The invention further discussed the possible involvement of AUR1 and AUR2 kinases in cancer and/or other signal transduction disorders, and the possible biol., diagnostic and/or therapeutic uses of these kinases. The AUR1 and AUR2 genes are mapped to chromosome 17p13.1 and 20q13.2 resp. Methods for treatment, diagnosis, and screening are provided for AUR1 and/or AUR2 related diseases or conditions characterized by an abnormal interaction between a AUR1 and/or AUR2 polypeptide and a AUR1 and/or AUR2 binding partner.

RE.CNT 182 THERE ARE 182 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 4 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN
AN 2002:636823 CAPLUS
DN 137:165497
TI Method and kit for assaying protein phosphorylation enzyme activity, and

antibody used for assay
IN Taji, Shingo; Tamai, Katsuyuki; Kobayashi, Toshiko
PA Medical and Biological Laboratories Co., Ltd., Japan
SO Jpn. Kokai Tokkyo Koho, 29 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2002236125	A	20020823	JP 2001-29774	20010206
PRAI	JP 2001-29774		20010206		

AB A method and a kit are provided for assaying a human Aurora2 protein phosphorylation enzyme activity using an antibody capable of specifically recognizing and binding with the substrate phosphorylated with human Aurora2 protein phosphorylation enzyme. An antibody used for assaying a human Aurora2 protein phosphorylation enzyme activity is also provided. A method is also provided for screening a compound which inhibits or promotes the human Aurora2 protein phosphorylation enzyme activity. The phosphorylation activity of human Aurora2 protein phosphorylation enzyme is assayed by immunol. measuring the phosphorylation of its substrate using an antibody capable of specifically recognizing and binding with the substrate phosphorylated with human Aurora 2 protein phosphorylation enzyme (e.g., human Lats2 protein).

L3 ANSWER 5 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2002:430919 CAPLUS

DN 137:227360

TI Comparison of two aquatic alphaviruses, salmon pancreas disease virus and sleeping disease virus, by using genome sequence analysis, monoclonal reactivity, and cross-infection

AU Weston, Jonathan; Villoing, Stephane; Bremont, Michel; Castric, Jeanette; Pfeiffer, Martin; Jewhurst, Victoria; McLoughlin, Marian; Roedseth, Odd Magne; Christie, Karen Elina; Koumans, Joseph; Todd, Daniel

CS Department of Veterinary Sciences, The Queen's University of Belfast, Belfast, BT4 3SD, UK

SO Journal of Virology (2002), 76(12), 6155-6163

CODEN: JOVIAM; ISSN: 0022-538X

PB American Society for Microbiology

DT Journal

LA English

AB Cell culture isolates of salmon pancreas disease virus (SPDV) of farmed Atlantic salmon and sleeping disease virus (SDV) of rainbow trout were compared. Excluding the poly(A) tracts, the genomic nucleotide sequences of SPDV and SDV RNAs include 11,919 and 11,900 nucleotides, resp. Phylogenetic anal. places SPDV and SDV between the New World viruses of Venezuelan equine encephalitis virus and Eastern equine encephalitis virus and the Old World viruses of Aura virus and Sindbis virus. When compared to each other, SPDV and SDV show 91.1% nucleotide sequence identity over their complete genomes, with 95 and 93.6% amino acid identities over their nonstructural and structural proteins, resp. Notable differences between the two viruses include a 24-nucleotide insertion in the C terminus of nsP3 protein of SPDV and amino acid sequence variation at the C termini of the capsid and E1 proteins. Exptl. infections of Atlantic salmon and rainbow trout with SPDV and SDV confirmed that the disease lesions induced by SPDV and SDV were similar in nature. Although infections with SPDV and SDV produced similar levels of histopathol. in rainbow trout, SDV induced significantly less severe lesions in salmon than did SPDV. Virus neutralization tests performed with sera from exptl. infected salmon indicated that SPDV and SDV belonged to the same serotype; however, antigenic variation was detected among SDV and geog. different SPDV isolates by using monoclonal antibodies. Although SPDV and SDV exhibit minor biol. differences, we conclude on the basis of the close genetic similarity that SPDV and SDV

are closely related isolates of the same virus species for which the name Salmonid alphavirus is proposed.

RE.CNT 55 THERE ARE 55 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 6 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2002:195040 CAPLUS

DN 137:92550

TI Expression of NOS-2, COX-2 and Th1/Th2 cytokines in migraine

AU Martelletti, Paolo; Zicari, Alessandra; Realacci, Massimo; Fiore, Giuseppe; De Filippis, Sergio; Stirparo, Giuseppe; Denora, Paola; Solimeo, Maria Donata; Rinaldi, Cristina; Morrone, Stefania; Giacobuzzo, Mario
CS Internal Medicine, Headache Centre, Sant'Andrea Hospital, 2nd School of Medicine, La Sapienza University of Rome, Rome, I-00189, Italy.

SO Journal of Headache and Pain (2001), 2(Suppl. 1), S51-S56

CODEN: JHPOAT; ISSN: 1129-2369

PB Springer-Verlag Italia Srl

DT Journal

LA English

AB Nitric oxide (NO) probably plays an important role in the pathogenesis of migraine without aura (MWA). As the activation of NO-ergic cascade has been shown to be closely linked to cyclooxygenase pathway and to cause some differences in peripheral blood lymphocyte populations, we investigated if the Th1/Th2 balance in peripheral blood of MWA patients was affected in comparison to controls. Twenty-six MWA patients and 10 healthy controls (C) were enrolled in this study. Blood samples were taken at baseline (T0) and during an induced migraine attack (T1). Total RNA from human peripheral blood lymphocytes (PBLs) was isolated and reverse-transcribed to prepare complementary DNA. COX-2, NOS-2 and β -actin were amplified using PCR. PBLs from patients and controls were stimulated with phorbol 12-myristate 13-acetate plus ionomycin in the presence of brefeldin A. Cells were surface-stained with fluorochrome-conjugated anti-CD3 and anti-CD8 monoclonal antibodies (mAbs) and intracellularly stained with fluorochrome-conjugated anti-IFN- γ or anti-IL-4 mAbs. The level of cytokine expression was analyzed by gating on the CD4+ lymphocyte subset. Th1 and Th2 type cytokines (IFN- γ , IL-2, IL-4) were directly assayed in serum by ELISA. Preliminary results indicate that NOS-2 was upregulated in MWA patients at basal time if compared to controls, whereas after NOD administration NOS-2 was significantly decreased. COX-2 was upregulated in MWA patients at basal time and it had an opposite trend after NOD administration. The homeostatic Th1/Th2 balance defined by the IFN- γ or IL-4 cytokine expression was unchanged in MWA patients in comparison to controls, and NOD administration did not affect that pattern. The cell activation machinery was not altered after mitogenic activation, as shown by CD69 expression level. Cytokine serum levels showed no significant changes in all studied situations. This study confirms the relevance of the NOS/COX system in MWA but, in contrast with previous studies, excludes their effect and activation on peripheral cytokine production. More sophisticated exptl. models are needed to investigate the ability of NOS/COX to activate migraine pain.

RE.CNT 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 7 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN

AN 1995:941486 CAPLUS

DN 124:25278

TI Analysis of the hemagglutination activity domains of the Venezuelan equine encephalomyelitis and eastern equine encephalomyelitis viruses

AU Razumov, I. A.; Khusainova, A. D.; Agapov, E. V.; Gajdamovich, S. Ya.; Pereboev, A. V.; Kolykhalov, A. A.; Netesov, S. V.; Loktev, V. B.

CS State Research Center Virology and Biotechnology "Vector", Institute Molecular Biology, Koltsovo, 633159, Russia

SO Intervirology (1995), Volume Date 1994, 37(6), 356-60

CODEN: IVRYAK; ISSN: 0300-5526

PB Karger
DT Journal
LA English

AB The hemagglutination (HA) domains of the Venezuelan equine encephalomyelitis (VEE) and the eastern equine encephalomyelitis (EEE) viruses providing the interaction of virions and red blood cells were studied with the use of a panel of 17 hemagglutination inhibition (HI) monoclonal antibodies (MAbs). A highly conserved domain (C domain) forming alphavirus-group-reactive MAbs inhibited in the E2 protein of the VEE and EEE viruses. These MAbs inhibited HA of the western equine encephalomyelitis, Semliki Forest, Sindbis, Getah, Aura, Chikungunya and Pixuna viruses. The involvement of amino acid residues 59 and 232 in the formation of the C region was demonstrated by sequencing the gene encoding the E2 protein of three escape variants of the VEE virus.

L3 ANSWER 8 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN

AN 1994:241981 CAPLUS

DN 120:241981

TI Molecular studies of alphavirus immunogenicity

AU Strauss, J. H.

CS California Inst. Tech., Pasadena, CA, USA

SO Report (1992), Order No. AD-A261546, 46 pp. Avail.: NTIS

From: Gov. Rep. Announce. Index (U. S.) 1993, 93(14), Abstr. No. 341,843

DT Report

LA English

AB The alphaviruses consist of a group of 26 closely related viruses. Many of these viruses can cause disease in man, characterized by encephalitis, polyarthrititis, fever or rash, depending upon the virus. In the 2.5 yr of research supported under this contract the authors have mapped antigenic epitopes in the structural glycoproteins of alphaviruses that lead to neutralization of virus infectivity upon reaction with an antibody, and have determined the sequence relationships of a number of Sindbis-like alphaviruses to one another and to other alphaviruses. The authors found that a domain of glycoprotein E2 of alphaviruses, between residues of 170 and 220, was an important region for binding of monoclonal antibodies that neutralize virus infectivity, making it critical importance for the immune response required for protection from infection by the virus.. In the determination of the relationships of alphaviruses to one another,

the authors have determined complete or partial sequences of 8 different alphavirus RNAs. These include Ockelbo virus, a virus causing epidemic polyarthrititis in northern Europe, strains of Sindbis virus from Africa, India, Australia and New Zealand arid Aura virus from South America...

=> s ((Aurora-A) or STK-15) and monoclonal

4145 AURORA

495 AURORAS

4226 AURORA

(AURORA OR AURORAS)

21285116 A

501 AURORA-A

(AURORA(W)A)

424 STK

31 STKS

452 STK

(STK OR STKS)

1750137 15

4 STK-15

(STK(W)15)

148722 MONOCLONAL

543 MONOCLONALS

148789 MONOCLONAL

(MONOCLONAL OR MONOCLONALS)

L4 12 ((AURORA-A) OR STK-15) AND MONOCLONAL

=> duplicate remove L4

PROCESSING COMPLETED FOR L4

L5 12 DUPLICATE REMOVE L4 (0 DUPLICATES REMOVED)

=> d L5 bib abs 1-12

L5 ANSWER 1 OF 12 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2007:488925 CAPLUS

DN 147:7139

TI Subcellular localization of the spindle proteins Aurora

A, Mad2, and BUBR1 assessed by immunohistochemistry

AU Burum-Auensen, Espen; De Angelis, Paula M.; Schjoelberg, Aasa R.; Kravik, Katherine L.; Aure, Marit; Clausen, Ole Petter F.

CS The Pathology Clinic, Rikshospitalet-Radiumhospitalet Medical Center, Faculty of Medicine, University of Oslo, Oslo, Norway

SO Journal of Histochemistry and Cytochemistry (2007), 55(5), 477-486

CODEN: JHCYAS; ISSN: 0022-1554

PB Histochemical Society, Inc.

DT Journal

LA English

AB The spindle checkpoint, the primary mechanism to ensure that two daughter cells receive the same amount of DNA, is compromised in many malignant tumors and has been implicated as a contributor to aneuploidy and carcinogenesis. The extent of expression and subcellular localization of the spindle proteins Aurora A, Mad2, and BUBR1 varies considerably in different immunohistochem. (IHC) reports from archival tumor tissues. Given the conflicting reports in the literature about the localization of these proteins, we examined the subcellular localization of Aurora kinase A, Mad2, and BUBR1 in normal and cancerous human tissues by IHC. In normal tissues, Aurora A was mainly localized to the nucleus when monoclonal or purified polyclonal antibodies were used, and Mad2 was localized to the nucleus, whereas BUBR1 was localized to the cytoplasm. In malignant tissues, Aurora A showed addnl. staining in the cytoplasm in the majority of tumors analyzed. Furthermore, BUBR1 was also localized to both the nucleus and cytoplasm in a significant fraction of tumors. Subcellular localization of Mad2 was similar in normal and malignant tissues. Thus, the validity of some earlier IHC studies of Aurora A, Mad2, and BUBR1 should be reconsidered, indicating that high-quality antibodies and a high-alkaline antigen-retrieval technique are required to achieve optimal results. We conclude that the subcellular localizations of these spindle proteins are different, although they have overlapping biol. functions, and that Aurora A and BUBR1 undergo a shift in the subcellular localization during malignant transformation.

RE.CNT 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 2 OF 12 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2006:1279840 CAPLUS

DN 146:45539

TI Preparation of aminopyridine derivatives as selective Aurora-A inhibitors for treatment of cancer

IN Kato, Tetsuya; Kawanishi, Nobuhiko; Mita, Takashi; Ohkubo, Mitsuru; Shimomura, Toshiyasu

PA Banyu Pharmaceutical Co., Ltd., Japan


SO PCT Int. Appl., 151pp.

CODEN: PIXXD2

DT Patent

LA Japanese

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2006129842	A1	20061207	WO 2006-JP311179	20060530
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	WO 2006046734	A2	20060504	WO 2005-JP19957	20051025
	WO 2006046734	A3	20060921		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	US 2006106029	A1	20060518	US 2005-258447	20051025
PRAI	JP 2005-161156	A	20050601		
	WO 2005-JP19957	A	20051025		
	JP 2004-315152	A	20041029		
	US 2005-692537P	P	20050621		
OS	MARPAT 146:45539				
GI					



AB The title compds. I [A1 is (RbjCRbj')_{m2}; A2 is (RaiCRai')_{m1}; A3 is (Y2Rc)_{n1}CO(Y3Rd)_{n2}R; m1 and m2 each is 1, 2, or 3; n1 and n2 each is 0 or 1; i is an integer of 1 to m1; j is an integer of 1 to m2; R is optionally substituted aryl, heteroaryl, or cycloalkyl; Rai and Rai' each is hydrogen, alkyl; Rbj and Rbj' each is hydrogen, alkyl; Rc, Rd, and R1 each is hydrogen, alkyl; X1 is CH, CX1a, N; X1a is (un)substituted alkyl; X2 is CH, N, etc.; X3 is CH, CX3a, N; X3a is (un)substituted alkyl; X4 is CH or N; Y1, Y2, and Y3 are the same or different and each is CH or N; Z1 and Z2 are the same or different and each is CH or N; and W is a 5-membered aromatic heterocycle, e.g., pyrazole or thiazole] are prepared. Thus, (5-bromothiazol-2-yl)-(6-(4-benzoylpiperazin-1-ylmethyl)pyridin-2-yl)amine

was prepared in a multistep process from 2-aminothiazole and 2,6-dichloropyridine. Compds. of this invention showed IC50 values of 0.36 nM to 110 nM against Aurora-A; they showed IC50 values of 47 nM to 28000 nM against Aurora-B.

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 3 OF 12 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2006:411981 CAPLUS

DN 144:450734

TI Preparation of novel aminopyridines having Aurora A selective inhibitory action

IN Ohkubo, Mitsuru; Kato, Tetsuya; Kawanishi, Nobuhiko; Mita, Takashi; Shimomura, Toshiyasu

PA Banyu Pharmaceutical Co., Ltd., Japan

SO PCT Int. Appl., 49 pp.


CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 3

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2006046735	A1	20060504	WO 2005-JP19958	20051025
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	US 2006106029	A1	20060518	US 2005-258447	20051025
	EP 1828165	A1	20070905	EP 2005-799006	20051025
	R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, HR				
PRAI	JP 2004-315152	A	20041029		
	JP 2005-161156	A	20050601		
	US 2005-692537P	P	20050621		
	WO 2005-JP19958	W	20051025		
OS	MARPAT 144:450734				
GI					



AB The title compds. I [R1, R2 = H, lower alkyl, or alternatively R1 and R2 are combined together to form CH2; R3 = halo; R4 = halo or Me substituted

with 1-3 halogen atoms; X = CH, N; W = (un)substituted thiazolyl, pyrazolyl, thiadiazolyl; with the proviso which are Aurora A selective inhibitors useful in combination therapy of cancer, were prepared E.g., a multi-step synthesis of I [R1, R2 = H; R3 = F; R4 = Cl; X = CH; W = 2-thiazolyl], starting from (6-bromopyridin-2-yl)methanol, was given (no characterization data provided for intermediates). The compds. I exhibit excellent Aurora A selective activity (biol. data were provided for exemplified compds. I). Pharmaceutical compns. comprising the compound I alone or in combination with other antitumor agent(s) were disclosed.

RE.CNT 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 4 OF 12 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2006:412038 CAPLUS

DN 144:450735

TI Preparation of novel aminopyridine derivatives having selective Aurora-A protein kinase inhibitory effect

IN Ohkubo, Mitsuru; Kato, Tetsuya; Kawanishi, Nobuhiko; Mita, Takashi; Shimomura, Toshiyasu

PA Banyu Pharmaceutical Co., Ltd., Japan

SO PCT Int. Appl., 148 pp.



CODEN: PIXXD2

DT Patent

LA Japanese

FAN.CNT 3

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2006046734	A2	20060504	WO 2005-JP19957	20051025
	WO 2006046734	A3	20060921		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	AU 2005297848	A1	20060504	AU 2005-297848	20051025
	US 2006106029	A1	20060518	US 2005-258447	20051025
	EP 1813609	A2	20070801	EP 2005-799135	20051025
	R:				
	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, HR				
	WO 2006129842	A1	20061207	WO 2006-JP311179	20060530
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	IN 2007DN03927	A	20070831	IN 2007-DN3927	20070525
PRAI	JP 2004-315152	A	20041029		
	JP 2005-161156	A	20050601		
	US 2005-692537P	P	20050621		



AB The title compds. (I) or pharmaceutically acceptable salts or ester thereof [wherein m1, m2 = 1, 2, 3; n1, n2 = 0, 1; i = an integer of from 1 to m1; j = an integer of from 1 to m2; R = (un)substituted aryl, heteroaryl or cycloalkyl; Rai, Rai', Rbj, Rbj', Rc, Rd, Re = H, lower alkyl; X1 = CH, CX1a, N; wherein X1a = (un)substituted lower alkyl; X2 = CH, N; X3 = CH, N, CX3a; wherein X3a = (un)substituted lower alkyl; X4 = CH, N; 1 or 2 of X1-X4 is N; Y1, Y2, Y3 = CH, N; Z1, Z2 = CH, N; W = a 5-membered aromatic heterocycle of formula Q including pyrazole or thiazole; wherein W1 = CH, N, NH, O, S; W2 = CH, CW2a, N, NW2b, O, S; wherein W2a, W2b = H, halo, cyano, C1-2 alkyl, C3-5 cycloalkyl, 1 or 2 halo-substituted C1-2 alkyl] are prepared These compds. are selective inhibitors of Aurora-A protein kinase over Aurora-B protein kinase and exhibit synergistic anticancer activity in combination with other anticancer agents. An anticancer agent containing the compound I, and the combined use of the above anticancer agent with another anticancer agent are also disclosed. Thus, a mixture of 2.70 g 6-chloromethyl-N-(thiazol-2-yl)pyridin-2-amine, 4.00 g 1-(3-chloro-2-fluorobenzoyl)piperazine, and 6.25 mL N,N-diisopropylethylamine, and 30 mL DMF was stirred at 90° for 2 h to give, after workup and silica gel chromatog., 6-[(4-(3-chloro-2-fluorobenzoyl)piperazin-1-yl)methyl]-N-thiazol-2-ylpyridin-2-amine (II; R = H). II (R = H) and II (R = 2-methyl-2H-tetrazol-5-yl) showed IC50 of 0.67 and 0.32 nM against Aurora-A protein kinase, resp., and 440 and 190 nM against Aurora-B protein kinase, resp. They showed IC50 of 11.00 and 0.21 µM against human cervical cancer cell (HeLa S3), resp., and also showed synergistic antiproliferative activity against HeLa S3 cells in combination with paclitaxel.

their preparation, pharmaceutical compositions, and use in therapy
 IN Dickson, John K., Jr.; Hodge, Carl Nicholas; Mendoza, Jose Serafin; Chen,
 Ke
 PA Amphora Discovery Corporation, USA
 SO PCT Int. Appl., 141 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2006020767	A2	20060223	WO 2005-US28549	20050811
	WO 2006020767	A3	20061109		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,				
	CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,				
	GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ,				
	LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA,				
	NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK,				
	SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU,				
	ZA, ZM, ZW				
	RW:				
	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,				
	IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ,				
	CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH,				
	GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,				
	KG, KZ, MD, RU, TJ, TM				
	AU 2005272815	A1	20060223	AU 2005-272815	20050811
	CA 2575466	A1	20060223	CA 2005-2575466	20050811
	US 2006052416	A1	20060309	US 2005-202927	20050811
	EP 1781287	A2	20070509	EP 2005-803385	20050811
	R:				
	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,				
	IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL,				
	BA, HR, MK, YU				
	IN 2007DN00668	A	20070803	IN 2007-DN668	20070124
PRAI	US 2004-601266P	P	20040813		
	US 2004-608834P	P	20040910		
	WO 2005-US28549	W	20050811		
OS	MARPAT 144:233067				
GI					

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The invention relates to 2-amidothiazole compds. of formula I, which are inhibitors of ATP-utilizing enzymes, such as synthetases, ligases, and kinases. In compds. I, R is OH, alkoxy, (un)substituted amino, (un)substituted cycloalkyl, (un)substituted aryl, or (un)substituted heteroaryl; L is a bond, carbonyl, -NHC(O)-, (un)substituted C1-4 alkylene, C1-4 alkylene-NHC(O)-, or C1-4 alkylene-C(O)-; W is selected from H, halo, (un)substituted alkyl, (un)substituted cycloalkyl, (un)substituted heterocyclyl, (un)substituted aryl, and (un)substituted heteroaryl; Q is (un)substituted alkyl, (un)substituted cycloalkyl, (un)substituted heterocyclyl, (un)substituted aryl, or (un)substituted heteroaryl; and Z is (un)substituted alkyl; with several provisos. The invention also relates to the preparation of I, pharmaceutical compns. comprising a therapeutically effective amount of a compound I, optionally one or more addnl. therapeutic agents, and at least one pharmaceutically acceptable vehicle, as well as to the use of the compns. for the treatment of conditions associated with ATP-utilizing enzymes. Addition of tert-Bu 3-aminopropanoate (β -alaninate) to N-Fmoc-isothiocyanate followed by deprotection gave thiourea II, which cyclized with 2-(bromoacetyl)benzofuran to give aminothiazole III. Amine III was acylated with thiophene-2-carbonyl chloride followed by ester cleavage and

amidation with nipecotamide (piperidine-3-carboxamide), resulting in the formation of amidothiazole IV. Some compds. of the invention express IC50 values of less than 30 μ M in cellular proliferation assays and some express EC50 values of less than 30 μ M in an assay for the induction of apoptosis in target cells.

L5 ANSWER 6 OF 12 CAPLUS COPYRIGHT 2007 ACS on STN
 AN 2006:99983 CAPLUS
 DN 144:184708
 TI Use of K-252a and kinase inhibitors for the prevention or treatment of HMGB1-associated pathologies
 IN Fumero, Silvano; Pilato, Francesco, P.; Barone, Domenico; Bertarione, Rava, Rossa, Luisa; Mainero, Valentina; Traversa, Silvio
 PA Creabilis Therapeutics S.p.A., Italy; Bio3research Srl
 SO PCT Int. Appl., 63 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2006010628	A1	20060202	WO 2005-EP8258	20050729
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	AU 2005266447	A1	20060202	AU 2005-266447	20050729
	CA 2575272	A1	20060202	CA 2005-2575272	20050729
	EP 1771178	A1	20070411	EP 2005-778429	20050729
	R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR			
PRAI	US 2004-591880P	P	20040729		
	US 2005-647007P	P	20050127		
	WO 2005-EP8258	W	20050729		

AB The present invention relates to the use of K-252a, a physiolo. active substance produced by microorganisms, and/or a kinase inhibitor and of its salts or synthetic and/or chemical modified derivs. for the prevention or treatment of HMGB1-associated pathologies. More particularly, the present invention relates to the use of K-252a for the prevention or treatment of restenosis.

RE.CNT 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 7 OF 12 CAPLUS COPYRIGHT 2007 ACS on STN
 AN 2006:1309859 CAPLUS
 DN 146:179238
 TI Phospho-regulation of human protein kinase Aurora-A: Analysis using anti-phospho-Thr288 monoclonal antibodies
 AU Ohashi, S.; Sakashita, G.; Ban, R.; Nagasawa, M.; Matsuzaki, H.; Murata, Y.; Taniguchi, H.; Shima, H.; Furukawa, K.; Urano, T.
 CS Department of Biochemistry II, Nagoya University Graduate School of Medicine, Showa-ku, Nagoya, Japan
 SO Oncogene (2006), 25(59), 7691-7702
 CODEN: ONCNES; ISSN: 0950-9232
 PB Nature Publishing Group
 DT Journal

LA English
 AB Mammalian Aurora-A is related to a serine/threonine protein kinase that was originally identified by its close homol. with Saccharomyces cerevisiae Ipllp and Drosophila melanogaster aurora that are key regulators in the orchestration of mitotic events. The protein level of Aurora-A, its peak kinase activity during mitosis, and its activation have been attributed to phosphorylation. Here we show that this enzyme is an arginine-directed kinase and define its substrate specificity. We also found that Thr288 within the activation loop is a critical residue for activating phosphorylation events in vitro and that it is spatiotemporally restricted to a brief window at mitosis on duplicated centrosomes and on spindle microtubules proximal to the poles in vivo. Immunodepletion assays indicated that an upstream kinase(s) of Aurora-A might exist in mammalian cells in addition to autophosphorylation. Furthermore, human activated Aurora-A forms complexes with the neg. regulator protein serine/threonine phosphatase type 1 (PP1) that was neg. phosphorylated on Thr320. Interestingly, phospho-specific Aurora-A monoclonal antibodies restrain Aurora-A kinase activity in vitro, providing further therapeutic avenues to explore.

RE.CNT 79 THERE ARE 79 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 8 OF 12 CAPLUS COPYRIGHT 2007 ACS on STN
 AN 2005:33225 CAPLUS
 DN 142:112460
 TI Monoclonal antibodies to fragment of human mitotic kinase Aurora-A phosphorylated at threonine 288, preparation, and use in cancer therapy
 IN Urano, Takeshi; Furukawa, Koichi
 PA Farma Design Inc., Japan
 SO Jpn. Kokai Tokkyo Koho, 16 pp.
 CODEN: JKXXAF
 DT Patent
 LA Japanese
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2005006532	A	20050113	JP 2003-172730	20030618
PRAI	JP 2003-172730		20030618		

AB This invention relates to antibodies, particularly, monoclonal antibodies, against human mitotic kinase Aurora-A (Aur-A) phosphorylated at threonine 288 (Thr-288), production in hybridoma, and use in treatment of diseases associated with Aur-A (over)expression, notably cancer. Monoclonal antibodies (mAbs) were raised against human Thr-288 phosphorylated Aur-A fragment. The mAbs were able to inhibit activation of Aur-A via phosphorylation of Thr-288.

L5 ANSWER 9 OF 12 CAPLUS COPYRIGHT 2007 ACS on STN
 AN 2004:1059119 CAPLUS
 DN 142:32932
 TI Combination therapy for cancer and other proliferative disorders
 IN Blatt, Lawrence M.; Seiwert, Scott D.; Ozes, Osman N.
 PA Intermune, Inc., USA
 SO PCT Int. Appl., 635 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004105684	A2	20041209	WO 2004-US15346	20040513

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,

GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
 LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI,
 NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,
 TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE,
 SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,
 SN, TD, TG

PRAI US 2003-471841P P 20030516
 US 2003-485474P P 20030708
 US 2003-511259P P 20031014
 US 2003-511280P P 20031014
 US 2003-511415P P 20031014
 US 2003-514173P P 20031024
 US 2004-561940P P 20040413

AB The invention provides methods of treating proliferative disorders, including angiogenesis-mediated disorders, cancer, and fibrotic disorders. In some embodiments, the methods involve administering a Type II interferon receptor agonist and a Type I interferon receptor agonist. In other embodiments, the methods involve administering a Type II interferon receptor agonist, a stress-activated protein kinase (SAPK) inhibitor, and a third therapeutic agent. In other embodiments, the methods involve administering a Type II interferon receptor agonist and a vascular endothelial growth factor (VEGF) antagonist. In other embodiments, the methods involve administering a VEGF antagonist and a SAPK inhibitor. The invention further provides methods of treating fibrotic disorders. In some embodiments, the methods involve administering a Type I interferon receptor agonist, a Type II interferon receptor agonist; and a tumor necrosis factor (TNF) antagonist. In other embodiments, the methods involve administering a Type II interferon receptor agonist and a TNF antagonist. In other embodiments, the methods involve administering pirfenidone or a pirfenidone analog and a TNF antagonist. In other embodiments, the methods involve administering a Type II interferon receptor agonist and a transforming growth factor- β (TGF- β) antagonist. In other embodiments, the methods involve administering a SAPK inhibitor alone or in combination with a Type II interferon receptor agonist. In other embodiments, the methods involve administering N-acetyl cysteine (NAC) and a SAPK inhibitor. In other embodiments, the methods involve administering NAC and a Type II interferon receptor agonist.

L5 ANSWER 10 OF 12 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2004:1079852 CAPLUS

DN 142:69142

TI Protein phosphatase CDC25B phosphopeptides, anti-phosphopeptide antibodies, and methods for cancer diagnosis and drug screening

IN Ducommun, Bernard; Monsarrat, Bernard; Prigent, Claude

PA Centre National de la Recherche Scientifique CNRS, Fr.

SO Fr. Demande, 32 pp.

CODEN: FRXXBL

DT Patent

LA French

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	FR 2856068	A1	20041217	FR 2003-7095	20030612
	FR 2856068	B1	20050819		
	CA 2528844	A1	20041223	CA 2004-2528844	20040608
	WO 2004111215	A1	20041223	WO 2004-FR1416	20040608
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,				

TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE,
 SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,
 SN, TD, TG

EP 1631665 A1 20060308 EP 2004-767282 20040608
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK

PRAI FR 2003-7095 A 20030612
 WO 2004-FR1416 W 20040608

AB A phosphopeptide derived from human protein phosphatase CDC25B, i.e.,
 TPVQNKRRRSpVTPPEEQQE, is disclosed. Also disclosed are polyclonal or
 monoclonal antibodies binding to this phosphopeptide. These
 antibodies may be used in diagnosis of breast cancer or in screening for
 antitumor agents. Thus, the site of phosphorylation of human CDC25B and
 its splice variants by protein kinase aurora A/STK5
 was determined

RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 11 OF 12 CAPLUS COPYRIGHT.2007 ACS on STN
 AN 2003:990980 CAPLUS
 DN 140:40888
 TI Monoclonal antibodies to Aurora A kinase and
 their use in the diagnosis and treatment of cancer
 IN Prigent, Claude; Martin, Anne
 PA Centre National De La Recherche Scientifique Cnrs, Fr.
 SO Fr. Demande, 33 pp.
 CODEN: FRXXBL
 DT Patent
 LA French
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	FR 2840905	A1	20031219	FR 2002-7212	20020612
	FR 2840905	B1	20060707		
	CA 2489214	A1	20031224	CA 2003-2489214	20030612
	WO 2003106500	A1	20031224	WO 2003-FR1772	20030612
	W:			AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW	
	RW:			GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG	
	AU 2003255671	A1	20031231	AU 2003-255671	20030612
	EP 1511771	A1	20050309	EP 2003-760023	20030612
	R:			AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK	
	JP 2006513135	T	20060420	JP 2004-513330	20030612
	US 2007117163	A1	20070524	US 2005-517645	20050210
PRAI	FR 2002-7212	A	20020612		
	WO 2003-FR1772	W	20030612		

AB The present invention has as an aim a monoclonal antibody
 directed against kinase aurora-A of the mammals, its
 process of obtaining, as its uses within the framework of the diagnosis or
 the forecast of cancers, and in pharmaceutical compns. within the
 framework of the treatment of cancers. Monoclonal antibodies
 have been raised against the Aurora A kinase for use
 in the diagnosis, prognosis, and treatment of cancer. The

monoclonal antibody 35C1 does not inhibit Aurora
A kinase.

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 12 OF 12 CAPLUS COPYRIGHT 2007 ACS on STN
AN 2002:958062 CAPLUS
DN 138:285769
TI Preparation and characterization of a human aurora-A
 kinase monoclonal antibody
AU Cremet, Jean Yves; Descamps, Simon; Verite, Frank; Martin, Ann; Prigent,
 Claude
CS Faculte de medecine, IFR 97 Genomique et Sante, CNRS - UMR 60611 Genetique
 et Developpement, Universite de Rennes 1, Rennes, 35043, Fr.
SO Molecular and Cellular Biochemistry (2003), 243(1&2), 123-131
 CODEN: MCBIB8; ISSN: 0300-8177
PB Kluwer Academic Publishers
DT Journal
LA English
AB We have developed monoclonal antibodies against the human
 aurora-A serine/threonine kinase. After immunization of
 a mouse, a fusion was performed to obtain hybridomas that were selected
 because they produced Ig pos. reacting against the protein used for
 immunization. We isolated one particular monoclonal that we
 named 35C1 using a series of selective assays. The first criteria of the
 screen for monoclonals was an Elisa (Enzyme Linked Immunosorbant
 Assay) assay performed in 96-well plates against the purified recombinant
 histidine-tagged aurora-A. The second was a pos.
 Western blot against the same recombinant protein. The third criteria was
 a pos. western blot against an HeLa cell extract, the selected
 monoclonal should detect only one protein migrating at 46 kDa
 (kiloDalton) on SDS (Sodium Dodecyl Sulfate)-polyacrylamide gel
 electrophoresis. Finally, the monoclonal had to bind to
 duplicated centrosomes and spindle poles in human MCF7 cultured cells by
 indirect immunofluorescence. At this stage several monoclonals
 were still pos. We then increased the selectivity by searching for
 antibodies that were able to cross-react with the mouse aurora-
 A kinase both by western blot and indirect immunofluorescence. We
 selected and cloned the 35C1 hybridoma to produce the antibody. Further
 characterization of the 35C1 antibody revealed that it was able to
 immunoppt. the kinase, that it did not inhibit the aurora-
 A kinase activity and consequently could be used to measure the
 aurora-A kinase activity in vivo after immunopptn.

RE.CNT 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> s (aurora-related kinase 1) or (hark1) or (breast-tumor-amplified kinase)

4145 AURORA
495 AURORAS
4226 AURORA
 (AURORA OR AURORAS)
1150462 RELATED
 1 RELATEDS
1150463 RELATED
 (RELATED OR RELATEDS)
299443 KINASE
57417 KINASES
308775 KINASE
 (KINASE OR KINASES)
9279975 1
 4 AURORA-RELATED KINASE 1
 (AURORA(W) RELATED(W) KINASE(W) 1)
 0 HARK1

79660 BREAST
 680 BREASTS
 79877 BREAST
 (BREAST OR BREASTS)
 421181 TUMOR
 161919 TUMORS
 472108 TUMOR
 (TUMOR OR TUMORS)
 80731 AMPLIFIED
 299443 KINASE
 57417 KINASES
 308775 KINASE
 (KINASE OR KINASES)
 5 BREAST-TUMOR-AMPLIFIED KINASE
 (BREAST(W)TUMOR(W)AMPLIFIED(W)KINASE)
 L6 9 (AURORA-RELATED KINASE 1) OR (HARK1) OR (BREAST-TUMOR-AMPLIFIED
 KINASE)

=> s 16 and monoclonal
 148722 MONOCLONAL
 543 MONOCLONALS
 148789 MONOCLONAL
 (MONOCLONAL OR MONOCLONALS)

L7 1 L6 AND MONOCLONAL

=> d 17 bib abs 1

L7 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2007 ACS on STN
 AN 2002:555652 CAPLUS
 DN 137:120714
 TI Human cDNA sequences and their encoded proteins and diagnostic and
 therapeutic uses
 IN Edinger, Shlomit; MacDougall, John R.; Millet, Isabelle; Ellerman, Karen;
 Stone, David J.; Gerlach, Valerie; Grosse, William M.; Alsobrook, John P.,
 II; Lepley, Denise M.; Rieger, Daniel; Burgess, Catherine E.; Casman,
 Stacie J.; Spytek, Kimberly A.; Boldog, Ferenc L.; Li, Li; Padigaru,
 Muralindhara; Mishra, Vishnu; Patturajan, Meera; Shenoy, Suresh; Rastelli,
 Luca; Tchernev, Velizar T.; Vernet, Corine A. M.; Zerhusen, Bryan D.;
 Malyankar, Uriel M.; Guo, Xiyojia; Miller, Charles E.; Gangolli, Esha A.
 PA Curagen Corporation, USA
 SO PCT Int. Appl., 353 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 175

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002057450	A2	20020725	WO 2001-US48922	20011129
	WO 2002057450	A3	20030717		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	AU 2002246696	A1	20020730	AU 2002-246696	20011129
	US 2004029116	A1	20040212	US 2002-87684	20020301
	US 7109000	B2	20060919		
	US 2004029222	A1	20040212	US 2002-218779	20020814
	AU 2005200106	A1	20050210	AU 2005-200106	20050112

	AU 2006201467	A1	20060504	AU 2006-201467	20060407
PRAI	US 2000-253834P	P	20001129		
	US 2000-250926P	P	20001130		
	US 2001-264180P	P	20010125		
	US 2001-313656P	P	20010820		
	US 2001-327456P	P	20011005		
	US 2001-995514	A	20011128		
	AU 2000-37360	A3	20000309		
	AU 2000-78680	A3	20001006		
	US 2001-274194P	P	20010308		
	WO 2001-US48922	W	20011129		

AB Disclosed herein are 12 cDNA sequences that encode novel human polypeptides that are members of the following protein families: transmembrane receptor UNC5H2-like, tyrosine phosphatase precursor-like, glomerular mesangial cell receptor protein tyrosine phosphatase precursor-like, Drosophila pecanex-like, Aurora-related kinase 1-like, 26S protease regulatory subunit 4-like, mitsugumin29-like, Wnt-15-like, Wnt-14-like, β -adrenergic receptor kinase-like, α -mannosidase-like, Clq-related factor-like, and plexin 1-like. Also disclosed are polypeptides encoded by these nucleic acid sequences, and antibodies, which immunospecifically-bind to the polypeptide, as well as derivs., variants, mutants, or fragments of the aforementioned polypeptide, polynucleotide, or antibody. The invention further discloses therapeutic, diagnostic and research methods for diagnosis, treatment, and prevention of disorders involving any one of these novel human nucleic acids and proteins.